



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# ENUMERATION OF PRIMES.

BY PROF. W. W. JOHNSON, ST. JOHN'S COLLEGE, ANNAPOLIS, MD.

MR. J. W. L. GLAISHER's enumeration of primes, referred to at page 9, Vol. II of the ANALYST, has been completed, and a "preliminary account" of the results appeared in the Proceedings of the Cambridge Philosophical Society, Vol. III.

In this paper, Mr. Glaisher gives tables of the form described below, for the first three millions of natural numbers comprised in Burckhardt's Tables of Divisors, and for the 7th, 8th and 9th millions which are covered by Dase's Tables.

Calling the hundred numbers between  $100n$  and  $100(n+1)$  a "century", the tables show, for each of the thirty groups of 100,000 in these six millions, the number of centuries which contain no prime, the number which contain a single prime, two primes, three, and so on; also the total number of primes in each group. These tables are given below, the ten numbers in each column being the numbers of primes in each of the ten groups of the million to which it refers, and the total at the foot, the number of primes in the entire million. The differences are appended to exhibit more strikingly the remarkable inequality which exists in the distribution of primes.

	1st million	2nd million	3rd million
1st group	9,593	7,216	6,874
2nd "	8,392 — 1,203	7,225 + 9	6,857 — 17
3rd "	8,013 — 379	7,081 — 144	6,849 — 8
4th "	7,863 — 150	7,103 + 22	6,791 — 58
5th "	7,678 — 185	7,028 — 75	6,770 — 21
6th "	7,560 — 118	6,973 — 55	6,809 + 39
7th "	7,445 — 115	7,015 + 42	6,765 — 44
8th "	7,408 — 37	6,932 — 83	6,716 — 49
9th "	7,323 — 85	6,957 + 25	6,746 + 30
10th "	7,224 — 99	6,903 — 54	6,708 — 38
	— 8	— 29	
Total	78,499	70,433	67,885

	7th million	8th million	9th million
1st group	6,397	6,369	6,250
2nd "	6,402 + 5	6,306 — 63	6,301 + 51
3rd "	6,425 + 23	6,348 + 42	6,283 — 18
4th "	6,337 — 88	6,299 — 49	6,285 + 2
5th "	6,347 + 10	6,301 + 2	6,245 — 40
6th "	6,402 + 55	6,305 + 4	6,245 + 81
7th "	6,338 — 64	6,347 + 42	6,326 — 45
8th "	6,375 + 37	6,245 — 102	6,281 + 19
9th "	6,411 + 36	6,364 + 119	6,299 — 79
10th "	6,365 — 46	6,274 — 90	6,220 + 50
	+ 4	— 24	
Total	<u>63,799</u>	<u>63,158</u>	<u>62,760</u>

Excepting the first two centuries, which contain 26 and 21 primes respectively, there are no centuries containing more than 17 primes; and excluding the first group of 100,000 numbers, there are no centuries containing more than 15 except two containing 17 primes, one between 700,000 and 800,000 and the other in the third million, and one containing 16 primes in the second million. Two centuries containing 15 primes occur in the eighth million, but none with more than 14 occur in the seventh or ninth. No centuries free from primes occur in the first million, and but one each in the second and third, but the seventh million contains six and the eighth and ninth four each, of such centuries.

Mr. Glaisher defers a comparison of his results with the logarithmic integral formula until after the completion of a sufficiently extended table of values of the integral.

PROPOSITION. BY L. W. MEECH, A. M.—If  $a, b, c$  denote three lines drawn from the angles of a plane triangle to the center of the inscribed circle whose radius is  $r$ , then will the reciprocals

$$\frac{1}{a}, \frac{1}{b}, \frac{1}{c}, \frac{1}{r}$$

be the four sides of a trapezium inscribed in a semicircle; the latter side coinciding with the diameter.